

Welcome to
MENTAL HEALTH MATTERS IN EARLY CHILDHOOD

THE TRAINING WILL BEGIN SHORTLY
While you're waiting...

CONNECT WITH US!

Icebreaker Question (answer in the chat)
 What is your favorite self-care activity or practice?

Survey & Certificate of Completion
 Available following the training.
 CEUs available for LCSWs, LMFTs, LPCCs, and LEPs

caltrin.org | twitter.com/cal_trin
 facebook.com/caltrin | linkedin/company/caltrin

1

SPANISH INTERPRETATION AVAILABLE!
INTERPRETACIÓN AL ESPAÑOL DISPONIBLE!

Today's training will be in both English and Spanish.
La capacitación de hoy será en español y inglés

Click the "Interpretation" icon in your toolbar, select "Spanish"

Seleccione el icono "Interpretation" de las opciones al debajo de su pantalla. Elige la opción "Spanish"

2

Hi, We're CalTrin!

Who we are

- The California Training Institute
- Funded by the State of California, Dept. of Social Services, Office of Child Abuse Prevention (OCAP) to support child abuse prevention through professional development and extended learning opportunities.
- Designed for staff of family strengthening and child abuse prevention organizations in California, including Family Resource Centers, Child Abuse Prevention Councils, community-based organizations, and other child and family serving systems.

What we offer

- Live webinars & small group training
- Virtual, self-paced courses
- Job aids & other resources

This training was made possible with funding from the California Department of Social Services, Office of Child Abuse Prevention. Any opinions, findings, conclusions, and/or recommendations expressed are those of the CEBC/CalTrin and do not necessarily reflect the views of the California Department of Social Services.

3

UPCOMING TRAININGS






mark your calendars!

Visit caltrin.org to view the full training calendar and the self-paced online training options


 09/29 Paternal Perinatal Mental Health: The Changing Face of New Fatherhood	 10/17 Parenting Traumatized Infants & Toddlers
 10/03 Addressing Implicit Bias	 10/20 The Art & Science of Father Engagement Programming
 10/10 Protective Factor: Concrete Support in Times of Need	 10/26 Rapid Response Team Collaboration

4

Housekeeping

DURING	DURING	AFTER
 Access the notetaking slides now! The link can be found in the chat.	 Review interactive features for today's session. Locate the controls on the toolbar at the bottom of your screen.	 Complete the survey to receive your Certificate of Attendance and CEUs.
 This presentation is being recorded.	<ul style="list-style-type: none"> • Questions in the Q&A • Discussion in the Chat 	 A follow-up email will be sent to all participants within 2 days.


5



Mental Health Matters in Early Childhood

Presenter: Meghan Lukasik, PhD
Yanan Guo, PsyD, BCBA

CHS is not responsible for the creation of content and any views expressed in its materials and programming.



6



Speakers SPOTLIGHT



YANAN GUO, PsyD, BCBA
Licensed Clinical Psychologist
 Developmental Evaluation Clinic
 Rady Children's Hospital-San Diego




MEGHAN LUKASIK, PHD
Manager & Licensed Psychologist
 Developmental Evaluation Clinic
 Rady Children's Hospital-San Diego

RCHSD is not responsible for the creation of content and any views expressed in its materials and programming.

7

AGENDA

- o Why mental health matter in early childhood
- o Signs and symptoms
- o History of mental health assessment
- o Trauma
- o Autism Spectrum Disorder (ASD)
- o Attention Deficit/Hyperactivity Disorder (ADHD)
- o Fetal Alcohol Spectrum Disorder (FASD)
- o Depression and anxiety
- o Resiliency and protective factors



8

The time is NOW

"Jim" age 26

- Incarcerated
- No family or friends
- History of violence
- Long history of drug and alcohol abuse



"Jimmy" age 10 months

- Removed from parents' care at birth
- Currently in 3rd resource/foster home
- Family history of domestic violence
- Born prenatally exposed to drugs and alcohol



9



10

Adult & Adolescent Mental Health By The Numbers

- 1 in 5 U.S. adults experience mental illness each year
- 1 in 20 U.S. adults experience serious mental illness each year
- 1 in 6 U.S. youth aged 6-17 experience a mental health disorder each year
- 50% of all lifetime mental illness begins by age 14, and 75% by age 24
- Suicide is the 2nd leading cause of death among people aged 10-34

NAMI: current

11

Early Childhood Mental Health Numbers

ADHD, behavior problems, anxiety, and depression are the most commonly diagnosed mental disorders in children ages 3-17


- 9.8% of children (approximately 6.0 million) have received an ADHD diagnosis.
- 9.4% of children (approximately 5.8 million) have diagnosed anxiety*
- 8.9% of children (approximately 5.5 million) have a diagnosed behavior problem
- 4.4% of children (approximately 2.7 million) have diagnosed depression*

1 in 6 U.S. children aged 2-8 years (17.4%) had a diagnosed mental, behavioral, or developmental disorder

*Depression and anxiety rates have increased over time


12

Signs of Wellness



- EAT
- SLEEP
- POOP
- REPEAT

13




Historical Perspective


1893: ICD-1	1949: ICD-6	1952: DSM-1	1968: DSM-II
1980: DSM-III	1987: DSM-III-R	1994: DSM-IV	1994: DC 0-3
2000: DSM-IV-TR	2005: DC 0-3R	2013: DSM 5	2017: DC 0-5
2022: DSM 5 TR			

14

Crosswalk : DSM5 - DC: 0-5 - ICD 10

- Neurodevelopmental Disorders
- Sensory Processing
- Anxiety Disorders
- Mood Disorders
- Obsessive-Compulsive and Related Disorders
- Sleep, Eating, and Crying Disorders
- Trauma, Stress, and Deprivation Disorders
- Relationship Disorder





Sleep, Eating, and Crying Disorders

Sleep Disorders


DC:0-5™	DSM-5	ICD-10	ICD-10 Code
Sleep Onset Disorder	Insomnia Disorder	Nonorganic Insomnia	F51.0

➔ Example

15

Trauma

- ▶ A thorough assessment starts with asking questions about stress and trauma exposure
- ▶ Types of Trauma
- ▶ Acute trauma
- ▶ Chronic trauma
- ▶ Complex trauma



16

Multiple Dimensions of Trauma

- ▶ The trauma
- ▶ Child's temperament or personality characteristics
- ▶ Caregiver's ability to help child cope and provide a sense of protection and safety



17




- Training
- Screening
- Protocols
- Payment
- Treatment



18

Adult Diseases Associated with Childhood Adversity & Impact on U.S. Health Care Costs



19

Pediatric ACEs and Related Life Events Screener (PEARLS)

www.acesaware.org

CHILD - To be completed by: Caregiver

At any point in time since your child was born, has your child seen or been present when the following experiences happened? Please include past and present experiences.

Please note, some questions have more than one part separated by "OR." If any part of the question is answered "Yes," then the answer to the entire question is "Yes."

PART 1:

- Has your child ever lived with a parent/caregiver who went to jail/prison?
- Do you think your child ever felt unsupported, unloved and/or unprotected?
- Has your child ever lived with a parent/caregiver who had mental health issues? (for example, depression, schizophrenia, bipolar disorder, PTSD, or an anxiety disorder)
- Has a parent/caregiver ever insulted, humiliated, or put down your child?
- Has the child's biological parent or any caregiver ever had, or currently has a problem with too much alcohol, street drugs or prescription medications use?

20

CHILD MEASURES

Name of measure	Area of assessment	Age range	Proprietary
Traumatic Events Screening Inventory - Child Self Report Revised (TESI-C SRR)	Traumatic events	8-17 years	no
Traumatic Events Screening Inventory - Parent Report Revised (TESI-PRR)	Traumatic events	4-7 years	no
Trauma History Questionnaire (THQ)	Traumatic events	7-17 years	no
Trauma Symptom Checklist for Children (TSCC)	Trauma related symptoms	8-16 years	yes
Trauma Symptom Checklist for Young Children (TSCYC)	Trauma related symptoms	3-12	yes

21

Trauma Informed Approach

- Understands the impact of trauma on behavior, development, and relationships Takes the child's developmental level into consideration
- Integrates this understanding into treatment planning
- Understands the provider's role in responding to child traumatic stress
- Reflects sensitively to the family



22

How is Early Childhood Trauma Unique?

- Profound sensory impact
 - Less able to anticipate danger or to know how to keep safe
 - Cannot express in words whether they feel afraid, overwhelmed or helpless
 - Early childhood trauma has been associated with reduced size of the brain cortex
 - Exclusive dependence on parents/caregivers for survival and protection—both physical and emotional


National Childhood Traumatic Stress Network (NCTSN) www.nctsn.org

23

Figure 8

THE IMPACT OF DEVELOPMENTAL EXPERIENCE

THE BALANCE BETWEEN ADVERSITY AND CONNECTEDNESS



The diagram is a 2x2 matrix with 'DEVELOPMENTAL RISK' on the vertical axis (Low at top, High at bottom) and 'GLOBAL HEALTH PROBLEMS' on the horizontal axis (Low on left, High on right). The vertical axis is labeled 'Adversity' (increasing downwards) and the horizontal axis is labeled 'Connectedness' (increasing rightwards). A blue dashed line runs from the top-left (Low Risk, Low Problems) to the bottom-right (High Risk, High Problems). A black dashed line runs from the top-right (Low Risk, High Problems) to the bottom-left (High Risk, Low Problems). A blue triangle is positioned in the bottom-left quadrant, pointing towards the origin.

First 2 months of life

With high connectedness and low adversity during development (blue dashed line), the balance of developmental risk is tipped in the direction of lower risk for mental, social, and physical health problems. In contrast, high adversity and minimal connectedness (black dashed line) increases developmental risk and the probability of significant problems in overall health.

24

Symptoms and Behaviors Associated with Exposure to Trauma

- Returning to behaviors shown at earlier ages
- Problems with toileting (bedwetting, soiling)
- Thumb sucking
- Fear of the dark
- Loss of language skills and acquired language
- Memory problems
- More immature behaviors
- Aimless motion, disorganized behaviors, and/or freezing
- Behavior changes
- Fear of being separated from parent/caregiver
- More clinging and dependent behaviors
- More aggressive behaviors
- More withdrawn behaviors showing little emotion
- More crying, whimpering, screaming, tantrums
- Unable to comfort self
- Difficulty falling asleep, night waking
- Less ability to tolerate frustration



National Childhood Traumatic Stress Network (NCTSN) www.nctsn.org

25

Effects of Trauma Exposure

- Neurological / Biological:**
 - Disruptions in biological / regulatory rhythms
 - Fundamental changes to structure and function of developing brain
 - Problems with movement and sensation
 - Somatic symptoms, increased medical problems.
- Self-Regulation:**
 - Challenges regulating affect, attention, action, and arousal
 - Difficulty knowing and describing their feelings and internal states.
- Attachment**
 - Challenges to basic trust and safety
 - Socially indiscriminate, lack of selectivity or preference
 - Attachment disturbances, insecure patterns of attachment
- Developmental:**
 - Regression - loss of previously acquired skills or disruption in developmental progress
 - Exacerbation of normative developmental fears

26

Effects of Trauma Exposure (Cont.)

- Social relatedness:**
 - Lack of consistent or discernable engagement vs. disengagement cues
 - Avoidance or indiscriminate patterns of social interaction
 - Difficulties with pro-social skills; sibling relational aggression
- Behavioral control**
 - Poor impulse control
 - Self-destructive or injurious behavior
 - Aggression
- Cognition:**
 - Difficulty focusing on and completing tasks, or planning for and anticipating future events.
 - Some exhibit learning difficulties and problems with language development.

27

The Impact of Trauma: Still Face Experiment

- <https://youtu.be/1eHGsFqK7So>



28

Autism Spectrum Disorder

A behaviorally described disorder affecting 1:36 children

Core features include difficulties With:

- Social Skills
- Verbal and Nonverbal Communication
- Repetitive behaviors /Restricted interests



29

Attention Deficit Hyperactivity Disorder (ADHD)

- A persistent pattern of **inattention** and/or **hyperactivity-impulsivity** that interferes with functioning or development:
- Several inattentive or hyperactive-impulsive symptoms present prior to age 12 years
- Several inattentive or hyperactive-impulsive symptoms present in two or more settings (e.g. at home, school or work; with friends or relatives; in other activities)
- Clear evidence that the symptoms interfere with life
- Symptoms are not better explained by another mental disorder or an event
- Different levels of severity



30

Fetal Alcohol Spectrum Disorders (FASD)


- Of all the substances people abuse ---including cocaine, heroin and marijuana--- alcohol produces the most serious neurobehavioral effects in the fetus
- No predictable correlation exists between the amount of alcohol exposure and the likelihood of development of an FASD
- During pregnancy, there is no safe time no safe amount, no safe type of alcohol



31


Criteria for FASD

- Growth defects
- Specific facial abnormalities
- Central Nervous System Abnormalities
 - Structural
 - Neurological
 - Functional



32

Depression



- Does the baby appear withdrawn, perhaps frequently staring into space?
- Does the baby's facial expression appear sad (infrequent smiling)?
- Is the baby expressing a vibrant range of emotions?
- Is it difficult to get your baby to engage with you socially?
- Is the baby quiet and subdued?
- Is there interest in toy play as expected for age?
- Is this behavior a change from the baby's usual presentation and temperament?


33

Anxiety



- Does your child have difficulty stopping themselves from worrying?
- Does your child have any fears?
- Does your child have to do things in the "right" order or position?
- Does your child ask for reassurance when it doesn't seem necessary?
- How does your child do when it is time to separate from you (preschool, daycare, babysitter, leaving the room)?
- **Need to consider frequency and intensity of behaviors to differentiate from developmentally appropriate reactions.**

34



Overlapping Symptoms of Childhood Disorders

- Trauma Reactions
- ASD
- ADHD
- FASD
- Depression
- ODD
- Anxiety
- Auditory Processing Disorder
- Sensory Processing Challenges
- OCD
- Gifted
- Motor Coordination Disorder

35

"Nathan's" Story



36

Changes in Cognition Over Time

	Differential Ability Scales 2 nd Edition	
	Age 3	Age 5
Verbal	92 (Average)	114 (Above Average)
Nonverbal Reasoning	100 (Average)	115 (Above Average)
Spatial	108 (Average)	123 (High)
General Cognitive Ability	100 (Average)	122 (High)

37

ADOS-2 and Diagnostic Impression Over Time

Age 2 Module 1	Age 3 Module 2	Age 5 Module 3
Just Met classification for autism spectrum	Met classification for autism spectrum	Non-spectrum
Adjustment Disorder primary	Adjustment Disorder primary	No diagnoses
Expressive Language Disorder	ASD Provisional	
Rule out Autism vs. Early signs of ADHD		

38

Strengthening Families Protective Factors Framework



Source <https://familynurturecenter.org/5-protective-factors/>

39

<https://youtu.be/Yn8j4XRxSck>



40

Questions?



41

Thanks for joining us!
WHAT'S NEXT?

- Survey and certificate in the chat now
- LCSWs, LPCCs, LMFTs, and LEPs – complete additional survey section to receive CEUs
- Follow-up email with resources within two days
- Watch your inbox for the next issue of *CalTrin Connect*



STAY CONNECTED FOR MORE FREE TRAINING & RESOURCES!



42
